

Single Colour Black
Dimensions: 110 x 150 mm
Cover Pg Substrate: 170 GSM ART CARD

Perforation lines
not to be print

INSTRUCTIONS FOR USE
Asenapine Sublingual Tablets

Read these Instructions for Use before you start using asenapine sublingual tablets and each time you get a refill. There may be new information. This leaflet does not take the place of talking to your doctor about your medical condition or your treatment.

IMPORTANT:

- For sublingual (under your tongue) use only
- Do not remove tablet until ready to administer.
- Use dry hands when handling tablet.

Directions for Taking your Asenapine Sublingual Tablets:

Step 1. Hold the blister as such that it faces the printed blister foil side and tear along the perforation marked on the blister to separate.




Figure A

Step 2. Peel the paper where “PEEL & PUSH” is given on the blister to expose foil.




Figure B

Step 3. Gently PUSH tablet through foil and take the tablet (see Figure C). Do not split, cut or crush the tablet.




Figure C

MSN LABORATORIES PRIVATE LIMITED		PACKAGING DEVELOPMENT	
Artwork Information		Specification for Printed Leaflet	
		Test	Specification
Brand Name	NA	Substrate	Maplitho Paper
Generic Name	Asenapine Sublingual Tablet	GSM	60 ± 5 % GSM
Open Dimension	110 X 150 ± 2 mm	Mode of supply	Binded Book Consists of 25 no's in each book
Folding Dimension	110 X 150 ± 2 mm	* Pharma Code	NA
Folding Type	Instruction in book form	Font Type	Arial Narrow, Arial Narrow Bold
Item Code	D03913-00	Font Size	10 pt
Supersede Code	NA	Perforation/Non-Perforation	NA
Country	USA	Developed by	Abdul Azeez
Customer	Breckenridge Pharmaceutical, Inc.	Reviewed by	N Raghu
Version	01	Colours	Black
Date & Time	22.07.2020 & 10.05		

Step 4. Place the **whole** tablet **under** tongue and allow it to dissolve completely (see Figure D).



Figure D

Do not chew or swallow the tablet.
Do not eat or drink for 10 minutes
(See Figure E).



Figure E

Storing asenapine sublingual tablets:

Store asenapine sublingual tablets at room temperature between 68°F to 77°F (20°C to 25°C).

These Instructions for Use have been approved by the U.S. Food and Drug Administration.

Manufactured by:
MSN Laboratories Private Limited
Telangana- 509 228, INDIA.

Distributed by:
Breckenridge Pharmaceutical, Inc.
Berlin, CT 06037

Revised: 07/2020

D03913-00

Perforation lines
not to be print